arXiv:1703.03826v2 [math.KT] 31 Mar 2017

ZERO-CYCLES WITH COEFFICIENTS FOR THE SECOND GENERALIZED SYMPLECTIC INVOLUTION VARIETY OF AN ALGEBRA OF INDEX 4

PATRICK K. MCFADDIN

Abstract. We compute the group of K_1-zero-cycles on the second generalized involution variety for an algebra of index 4 with symplectic involution. Our method utilizes the framework of Chernousov and Merkurjev for computing K_1-zero-cycles in terms of R-equivalence classes of prescribed algebraic groups. This gives a computation of K_1-zero-cycles for some homogeneous varieties of type C_n.

1. Introduction

The formalism of "Chow groups with coefficients," as developed by Markus Rost [Ros], has become a useful tool for investigating intersection-theoretic questions in arithmetic and algebraic geometry. This theory has bolstered the study of algebraic cycles by providing a broader context and more desirable functoriality properties than those found for classical Chow groups. Given a variety X defined over a field F, there is a complex [Qui, Ros]

$$\cdots \to \coprod_{x\in X_{(p+1)}} K_{p+q+1}(F(x)) \to \coprod_{x\in X_{(p)}} K_{p+q}(F(x)) \to \coprod_{x\in X_{(p-1)}} K_{p+q-1}(F(x)) \to \cdots$$

where $X_{(i)}$ is the set of points of X of dimension i, K_i are Quillen K-groups, and the differentials are given component-wise by residue and corestriction homomorphisms. The p^{th} homology group of this complex, i.e., homology at the middle term above, is denoted $A_p(X, K_q)$, and classical Chow groups in the sense of [Ful] are recovered via the identification $\mathrm{CH}_p(X) = A_p(X, K_{-p})$. The study of these K-cohomology groups for homogenous varieties has seen many useful applications to Galois cohomology, and significant results include the Merkurjev-Suslin Theorem [MS82] and the Milnor and Bloch-Kato Conjectures [Voe]. Unfortunately, general descriptions of these groups remain elusive, and computations are done in various cases, described below.

Algebraic cycles have also found a great deal of utility in the study of algebraic groups. In [CM01], Chernousov and Merkurjev show that for certain groups G and appropriately chosen homogenous varieties X for G, the group $A_0(X, K_1)$ encodes arithmetic properties of G that can be used to investigate rationality by means of R-equivalence. These results allow one the use of algebraic K-theory and K-cohomology in the analysis of such questions. In the present paper, we focus on the reverse

2010 *Mathematics Subject Classification.* Primary 14C25, Secondary 16K20.

implication. Our main result does not shed light on the rationality problem for the symplectic groups (shown to be rational in [Weil]), but nevertheless crucially utilizes the Chernousov-Merkurjev framework to identify K-cohomology groups.

Computing groups of zero-cycles (with coefficients) on homogeneous varieties has been taken up by a number of authors. The groups of K_0-zero-cycles, i.e., Chow groups $\mathrm{CH}_0(X)$, were computed for e.g. some (generalized) Severi-Brauer varieties, (generalized) involution varieties, Severi-Brauer flag varieties, and varieties of exceptional type in [CM06, Karp, Kra, Mer95, Pan84, PSZ, Swa].

For K_1-zero-cycles, many cases remain unknown. Let G be a simply connected group of inner type A_n, so that $G = \mathbf{SL}_1(A)$ for a central simple algebra A over F. For any $1 \leq k \leq \deg(A)$, the (generalized) Severi-Brauer varieties $X = \mathrm{SB}_k(A)$ associated to A are homogeneous for G and their K_1-zero-cycles have been computed in few cases. For X the Severi-Brauer variety of A (i.e., $k = 1$), Merkurjev and Suslin have shown that $A_0(X, K_1) \simeq K_1(A)$ [MS92]. For X the second generalized Severi-Brauer variety (i.e., $k = 2$) of an algebra A of index 4, it was shown by the author that $A_0(X, K_1) \simeq \{(x, \alpha) \in K_1(A) \times F^\times \mid \mathrm{Nrd}_A(x) = \alpha^2\}$ [McF].

For simply connected groups of types B_n and D_n, Chernousov and Merkurjev provide a description of the K_1-zero-cycles on the involution varieties associated to quadratic spaces and their twisted forms. Indeed, it was shown in [CM01] that for X the projective quadric defined by a quadratic form q on a vector space V, there is a canonical isomorphism $A_0(X, K_1) \simeq \Gamma^+(V, q)/R\mathrm{Spin}(V, q)$. If X is the involution variety corresponding to an algebra A of index no greater than 2 with quadratic pair (σ, f), there is an isomorphism $A_0(X, K_1) \simeq \Gamma^+(A, \sigma, f)/R\mathrm{Spin}(A, \sigma, f)$. Note that this work also recovers a theorem of Rost showing that $\overline{A}_0(X, K_1) = 0$ for X the projective quadric of a Pfister form [Ros88], used in the proof of the Bloch-Kato Conjecture [Voe].

In the present paper, we treat type C_n. Such groups are given by symplectic groups $G = \mathbf{Sp}(A, \sigma)$ for a central simple algebra of degree $2n$ with symplectic involution. The second generalized involution variety $\mathrm{IV}_2(A, \sigma)$ is homogeneous for G and we compute its K_1-zero-cycles for A of index 4. Our main result is the following:

Theorem (4.6)**.** *Let (A, σ) be a central simple algebra over F of index 4 with symplectic involution. Let $X = \mathrm{IV}_2(A, \sigma)$ be the corresponding second generalized involution variety. There is a canonical group isomorphism $A_0(X, K_1) \simeq F^\times$. Moreover, $\overline{A}_0(X, K_1) = 0$.*

Organization. In Section 2 we define a slew of homogeneous varieties and recall some structure results. We then define cycle modules and K-cohomology groups. In Section 3 we present the main algebraic groups of interest: spin, Clifford, (general) symplectic. We also recall some exceptional identifications of groups of type B_2 and C_2 and their algebraic invariants. Lastly, we define R-triviality and the index of an algebraic group. In Section 4 we present the framework of Chernousov and Merkurjev describing the group of K_1-zero-cycles in terms of R-equivalence classes. We provide our main result for symplectic involution varieties of algebras of degree 4. To obtain

our result for algebras of index 4, we analyze the fibers of two projection morphisms to relate cycles on certain involution and Severi-Brauer varieties.

Acknowledgements. We would like to thank Danny Krashen for helpful comments and suggestions.

2. Preliminaries

Throughout, F denotes an infinite perfect field of arbitrary characteristic. If L/F is a field extension and A is an F-algebra, we write $A_L := A \otimes_F L$. By a *scheme* we will mean a separated scheme of finite type over the field F. A variety is an integral scheme. If X and Y are F-varieties, we write $X_Y := X \times_{\operatorname{Spec} F} Y$.

A *central simple algebra* over F is a finite-dimensional F-algebra with no two-sided ideals other than (0) and (1) and whose center is precisely F. Recall that the dimension of a central simple algebra A is a square, and we define the *degree* of A to be $\deg(A) = \sqrt{\dim A}$. One may write $A = M_n(D)$ for a division algebra D, unique up to isomorphism, and we define the *index* of A to be $\operatorname{ind}(A) = \deg(D)$. The *period* (or exponent) of an algebra A is its order in the Brauer group of F.

An *involution* σ on a central simple algebra A is an anti-automorphism satisfying $\sigma \circ \sigma = \mathrm{id}_A$. An involution is *of the first kind* if $\sigma|_F = \mathrm{id}_F$. Upon extending scalars to a separable closure, involutions become adjoint to bilinear forms. Involutions which are adjoint to a skew-symmetric bilinear form (respectively, symmetric bilinear form) over a separable closure of F are *symplectic* (respectively, *orthogonal*). Note that an algebra with symplectic involution necessarily has even degree [KMRT, Prop 2.6]. For an algebra with involution of the first kind (A, σ), the set of *symmetrized elements* is

$$\operatorname{Symd}(A, \sigma) = \{a + \sigma(a) \mid a \in A\}.$$

2.1. Severi-Brauer and Involution Varieties.

See [Bla, KMRT, Kra, Tao]. Throughout this section, let A be a central simple algebra of degree n. Involutions on A are assumed to be of the first kind, either orthogonal or symplectic.

Definition 2.1. For a (right or left) ideal $I \subset A$, the *reduced dimension* of I is

$$\operatorname{rdim}(I) = \frac{\dim_F I}{n}.$$

Definition 2.2. For any integer $1 \leq k \leq n$, the k^{th} *generalized Severi-Brauer variety* $\mathrm{SB}_k(A)$ of A is the variety of right ideals of dimension nk (or reduced dimension k) in A.

Such a varietiy is a twisted form of the Grassmannian $\mathrm{Gr}(k, n)$ of k-dimensional subspaces of an n-dimensional vector space. The variety $\mathrm{SB}_1(A) = \mathrm{SB}(A)$ is the usual Severi-Brauer variety of A, which is a twisted form of projective space $\mathbb{P}^{n-1} = \mathrm{Gr}(1, n)$. For a field extension L/F, the variety $\mathrm{SB}_k(A)$ has an L-rational point if and only if $\operatorname{ind}(A_L) \mid k$ [KMRT, Prop. 1.17].

Definition 2.3. Let (A, σ) be a central simple algebra with involution σ (of the first kind, either orthogonal or symplectic). A right ideal I is *σ-isotropic* if $\sigma(I) \cdot I = (0)$. For any integer $1 \leq k \leq n$, the k^{th} *generalized involution variety* $\mathrm{IV}_k(A, \sigma)$ is the variety of σ-isotropic right ideals of reduced dimension k [Kra, §8].

Forgetting the isotropy condition defines an embedding $\mathrm{IV}_k(A, \sigma) \hookrightarrow \mathrm{SB}_k(A)$. In the case where $k = 1$ and σ is orthogonal, the above embedding realizes $\mathrm{IV}_1(A, \sigma)$ as a twisted projective quadric. This variety was first defined and studied in [Tao].

In Section 4, we make use of a homogeneous variety which interpolates between Severi-Brauer and involution varieties. We first provide the notion of a twisted flag variety to provide context.

Definition 2.4. For any collection of integers $1 \leq n_1 < n_2 < \cdots < n_k \leq n$, define the *Severi-Brauer flag variety* $V_{n_1,\ldots,n_k}(A)$ to be the variety parametrizing flags of right ideas $I_1 \subset \cdots \subset I_k \subset A$, where the reduced dimension of I_i is n_i [Kra].

Just as in the case of involution varieties, one generalizes the notion of flag variety to include the data of an involution. For our purposes, it will suffice to define these varieties using flags of length 2.

Definition 2.5. Let (A, σ) be an algebra with involution. For any pair of integers $1 \leq p < q \leq n$, define the *mixed involution flag variety* $V_{\overline{p},q}(A, \sigma)$ (resp. $V_{p,\overline{q}}(A, \sigma)$) to be the variety parametrizing flags $J \subset I$ where J has reduced dimension p, I has reduced dimension q, and J is σ-isotropic (resp. I is σ-isotropic).

There are natural projection maps $V_{\overline{p},q}(A, \sigma) \to \mathrm{IV}_p(A, \sigma)$ and $V_{\overline{p},q}(A, \sigma) \to \mathrm{SB}_q(A)$, defined by $J \subset I \mapsto J$ and $J \subset I \mapsto I$, respectively. Likewise for $V_{p,\overline{q}}(A, \sigma)$.

Definition 2.6. Let I be a right ideal of an algebra A. The *annihilator* of I is the left ideal of A given by $\mathrm{ann}(I) = \{x \in A \mid xI = (0)\}$. If σ is an involution on A, a right ideal $I \subset A$ is *σ-regular* if $I \cap \mathrm{ann}(\sigma(I)) = (0)$. The scheme $V_k(A)_{\mathrm{reg}}$ consisting of σ-regular right of ideals of A of reduced dimension k is an open subscheme of $V_k(A)$, so that A contains a σ-regular right ideal if and only if $\mathrm{ind}(A) \mid k$ [Kra, §8].

Definition 2.7. Let (A, σ) be an algebra with involution and I a right ideal of A. For any $1 \leq k \leq \mathrm{rdim}(I)$, let $\mathrm{IV}_k(I, \sigma)$ be subvariety of $\mathrm{IV}_k(A, \sigma)$ consisting of σ-isotropic varieties $J \subset I$ of reduced dimension k. [Kra, Def. 8.2]

Theorem 2.8 ([Kra], Thm. 8.3)**.** *Supppose (A, σ) is an algebra with involution. Let $I \subset A$ be a regular right ideal of reduced dimension k. Then there exists a degree k algebra D, Brauer-equivalent to A, with involution τ of the same type as σ and such that for any $1 \leq \ell \leq k$, we have*

$$\mathrm{IV}_\ell(I, \sigma) = \mathrm{IV}_\ell(D, \tau).$$

For algebras of degree 4, exceptional identifications allow us to recognize additional structure (see Section 3). A geometric consequence is given by the following result of Krashen.

Proposition 2.9 ([Kra], Prop. 8.10)**.** *Suppose A is a degree 4 central simple algebra with symplectic involution σ over a field F with* $\mathrm{char}(F) \neq 2$*. Then* $\mathrm{IV}_2(A, \sigma)$ *is isomorphic to a quadric hypersurface in* $\mathbb{P}^4$*.*

We explicitly identify this quadric hypersurface in terms of σ-symmetrized elements in A below (see Remark 3.5).

2.2. **Cycle Modules and Homology.** Cycle modules were introduced by M. Rost in [Ros] and good references are [EKM, GMS].

Definition 2.10. A *cycle module* M over F is a function assigning to every field extension L/F a graded abelian group $M(L) = M_*(L)$, which is a graded module over the Milnor K-theory ring $K^M_*(F)$ satisfying certain data and compatibility axioms. This data includes

(1) For each field homomorphism $L \to E$ over F, there is a degree 0 homomorphism $r_{E/L} : M(L) \to M(E)$ called *restriction*.
(2) For each field homomorphism $L \to E$ over F, there is a degree 0 homomorphism $c_{E/L} : M(E) \to M(L)$ called *corestriction* (or norm).
(3) For each extension L/F and each (rank 1) discrete valuation v on L, there is a degree -1 homomorphism $\partial_v : M(L) \to M(\kappa(v))$ called the *residue homomorphism*, where $\kappa(v)$ is the residue field of v.

These homomorphisms are compatible with the corresponding maps in Milnor K-theory (see D1-D4, R1a-R3e of [Ros, Def. 1.1]). For an F-scheme X, we say that a cycle module M over F is a cycle module on X if it further satisfies properties FD and C of [Ros, Def. 2.1] concerning those extension of F which arise as residue fields of points of X.

Let X be an F-variety and M a cycle module on X. Utilizing this axiomatic framework and setting

$$C_p(X, M_q) = \coprod_{x \in X_{(p)}} M_{p+q}(F(x)),$$

one obtains a complex

$$\cdots \to C_{p+1}(X, M_q) \xrightarrow{d_X} C_p(X, M_q) \xrightarrow{d_X} C_{p-1}(X, M_q) \to \cdots$$

with differentials d_X induced by the homomorphisms ∂_v associated to discrete valuations of subvarieties of codimension 1. This is often referred to as the *Rost complex* [EKM] or *Gersten complex* [Pan03]. We denote the homology group at the middle term by $A_p(X, M_q)$. Our main focus will be the group

$$A_0(X, K_1) = \mathrm{coker}\left(\coprod_{x \in X_{(1)}} K_2(F(x)) \xrightarrow{d_X} \coprod_{x \in X_{(0)}} K_1(F(x))\right)$$

of K_1-zero-cycles, where K_* is the cycle module given by Quillen or Milnor K-theory, as these coincide in low degree. References for Quillen K-theory include [Sri, Wei]

and for Milnor K-theory see [GS06]. The group of K_1-zero-cycles is therefore given by equivalences classes of expressions $\sum(\alpha, x)$, where x is a closed point on X and α and element of $F(x)^\times = K_1(F(x))$. Note that the group $A_0(X, K_1)$ coincides with the cohomology group $H^d(X, \mathcal{K}_{d+1})$, where $d = \dim(X)$ and $\mathcal{K}_i$ is the Zariski sheaf associated to the Quillen K-theory presheaf $U \mapsto K_i(U)$.

Definition 2.11. Let X be a complete F-variety. For $p \geq 0$ define the *norm homomorphism* $N^p : A_0(X, K_p) \to K_p(F)$ by

$$N^p\left(\sum(\alpha_x, x)\right) = \sum c_{F(x)/F}(\alpha_x),$$

where $c_{F(x)/F}$ is the corestriction. The kernel of N^p will be denoted $\overline{A}_0(X, K_p)$. In low degree (e.g., for $p = 0, 1$) these norm homomorphisms coincide with those defined in terms of Milnor K-theory [CM01, §1.5]. For the remainder, we denote the norm map N^0 by $\deg : \mathrm{CH}_0(X) \to \mathbb{Z}$, noting that $A_0(X, K_0) = \mathrm{CH}_0(X)$ and $K_0(F) = \mathbb{Z}$, and refer to it as the *degree homomorphism* [CM06]. Its image is equal to $\mathrm{ind}(X)\mathbb{Z}$.

3. Algebraic Groups

Let A be a central simple algebra over F, and define the *general linear group* $\mathbf{GL}_1(A)$ of invertible elements in A by $\mathbf{GL}_1(A)(R) = A_R^\times$ for any commutative F-algebra R. We write $\mathbb{G}_m = \mathbf{GL}_1(F)$. We will be interested in certain subgroups of $\mathbf{GL}_1(A)$ which reflect information about involutions on A.

Let (A, σ) be an algebra with symplectic involution. For any commutative F-algebra R, let

$$\mathbf{GSp}(A, \sigma)(R) = \{a \in A_R^\times \mid a \cdot \sigma_R(a) \in R^\times\}$$

$$\mathbf{Sp}(A, \sigma)(R) = \{a \in A_R^\times \mid a \cdot \sigma_R(a) = 1\}.$$

These are (non-respectively) the *symplectic* and *general symplectic* groups of (A, σ). In the above definition, $\sigma_R : A_R \to A_R$ is the involution obtained by extending scalars to R. Notice that we have a group homomorphism $\mu : \mathbf{GSp}(A, \sigma) \to \mathbb{G}_m$ defined by $\mu_R = a \cdot \sigma_R(a)$ on R-points. We call this the *multiplier map* (associated to σ) which clearly has kernel $\mathbf{Sp}(A, \sigma)$.

Now let (V, q) be a non-degenerate quadratic space and define the *Clifford algebra* $C(V, q)$ to be the quotient of the tensor algebra $T(V) = \bigoplus_i V^{\otimes i}$ by the ideal $I(q)$ generated by all elements of the form $v \otimes v - q(v) \cdot 1$ with $v \in V$. The $\mathbb{N}$-grading on $T(V)$ induces a $\mathbb{Z}/2$-grading on $C(V, q) = C_0(V, q) \oplus C_1(V, q)$, and we call $C_0(V, q)$ the *even Clifford algebra*. Define the *even Clifford group* by

$$\mathbf{\Gamma}^+(V, q)(R) = \{g \in C_0(V, q)_R^\times \mid g \cdot V_R \cdot g^{-1} = V_R\}.$$

The identity map on V extends to an involution on $T(V)$ which fixes the ideal $I(q)$. The induced involution on $C(V, q)$ restricts to $C_0(V, q)$ and we denote the resulting involution by τ, and call it the *canonical involution* on $C_0(V, q)$. The multiplier map

associated to τ defines a homomorphism $\mathrm{Sn} : \mathbf{\Gamma}^+(V,q) \to \mathbb{G}_m$ via $x \mapsto x \cdot \tau(x)$, called the *spinor norm* and its kernel is denoted

$$\mathbf{Spin}(V,q) = \{g \in \mathbf{\Gamma}^+(V,q)(R) \mid \ g \cdot \tau(g) = 1\}.$$

Remark 3.1. We follow the same notational convention as [CM01], utilizing $\mathrm{GSp}(A,\sigma)$, $\mathrm{Sp}(A,\sigma)$, $\Gamma^+(V,q)$, and $\mathrm{Spin}(V,q)$ to denote the groups of F-points of the corresponding algebraic groups $\mathbf{GSp}(A,\sigma), \mathbf{Sp}(A,\sigma), \mathbf{\Gamma}^+(V,q)$, and $\mathbf{Spin}(V,q)$.

Exceptional Identifications. Here we recall some facts given in [KMRT, §15.C]. Let B_2 be the groupoid of *oriented* quadratic spaces [KMRT, Def. 12.40] of dimension 5, where the morphisms are isometries which preserve the orientation. Let C_2 be the groupoid of central simple algebras of degree 4 with symplectic involution, where the morphisms are the F-algebra isomorphisms which preserve the involutions.

Define $\mathbf{C} : \mathsf{B}_2 \to \mathsf{C}_2$ via $(V,q,\zeta) \mapsto (C_0(V,q),\tau)$, where τ is the canonical involution on the even Clifford algebra $C_0(V,q)$.

The map in the other direction requires further development. Let (A,σ) be a central simple algebra of degree 4 with symplectic involution and consider the collection $\mathrm{Symd}(A,\sigma)$ of σ-symmetrized elements in A. The reduced characteristic polynomial of an element $a \in \mathrm{Symd}(A,\sigma)$ is a square and takes the form

$$\mathrm{Prd}_{A,a}(x) = \left(x^2 - \mathrm{Trp}_\sigma(a)x + \mathrm{Nrp}_\sigma(a)\right)^2.$$

Here, Trp_σ is the *Pfaffian trace* and is a linear form on $\mathrm{Symd}(A,\sigma)$; Nrp_σ is the *Pfaffian norm* and is a quadratic form on $\mathrm{Symd}(A,\sigma)$. Notice that if $\mathrm{Trp}_\sigma(a) = 0$, then $0 = a^2 + \mathrm{Nrp}_\sigma(a)$ since a satisfies its own characteristic polynomial.

Let $\mathrm{Symd}(A,\sigma)^0 = \{a \in \mathrm{Symd}(A,\sigma) \mid \mathrm{Trp}_\sigma(a) = 0\}$. Then $-\mathrm{Nrp}_\sigma(a) = a^2$ defines a quadratic form on $\mathrm{Symd}(A,\sigma)^0$ which we will denote by s_σ. We have thus associated a quadratic space (of dimension 5) to (A,σ). Lastly, there is a unique orientation η on this quadratic space, and we refer the reader to loc. cit. Define $\mathbf{S} : \mathsf{C}_2 \to \mathsf{B}_2$ via $(A,\sigma) \mapsto (\mathrm{Symd}(A,\sigma)^0, s_\sigma, \eta)$.

Proposition 3.2 ([KMRT], Prop. 15.16)**.** *The functors $\mathbf{C} : \mathsf{B}_2 \to \mathsf{C}_2$ and $\mathbf{S} : \mathsf{C}_2 \to \mathsf{B}_2$ define an equivalence of groupoids.*

Proposition 3.3 ([KMRT], Prop. 15.18)**.** *Suppose that $(A,\sigma) \in \mathsf{C}_2$ and $(V,q,\zeta) \in \mathsf{B}_2$ correspond to one another under the above groupoid equivalence. Then the special Clifford group and general symplectic group coincide: $\Gamma^+(V,q) = \mathrm{GSp}(A,\sigma)$. The spin group and symplectic group coincide: $\mathrm{Spin}(V,q) = \mathrm{Sp}(A,\sigma)$.*

The following proposition is in some sense an algebraic restatement of of Proposition 2.9, although there is no requirement that $\mathrm{char}(F) \neq 2$.

Proposition 3.4 ([KMRT], Prop. 15.20)**.** *(1). For every right ideal $I \subset A$ of reduced dimension 2, the intersection $I_\sigma := I \cap \mathrm{Symd}(A,\sigma)$ is a 1-dimensional subspace of $\mathrm{Symd}(A,\sigma)$ which is isotropic for the quadratic form Nrp_σ. If $\sigma(I) \cdot I = 0$ (i.e. if I is σ-isotropic), then $I_\sigma \subset \mathrm{Symd}(A,\sigma)^0$ and is thus s_σ-isotropic.*

(2). For every nonzero vector $x \in \mathrm{Symd}(A,\sigma)$ with $\mathrm{Nrp}_\sigma(x) = 0$, the right ideal xA has reduced dimension 2. Moreover, if $x \in \mathrm{Symd}(A,\sigma)^0$ then xA is σ-isotropic.

Remark 3.5. This proposition gives mutually inverse maps $I \mapsto I \cap \mathrm{Symd}(A,\sigma)$ and $xF \to xA$ between the sets of right ideals of A of reduced dimension 2 and Nrp_σ-isotropic subspaces of $\mathrm{Symd}(A,\sigma)$. Geometrically, this defines an isomorphism

$$\mathrm{SB}_2(A) \xrightarrow{\sim} \{\mathrm{Nrp}_\sigma = 0\} \subset \mathbb{P}(\mathrm{Symd}(A,\sigma)) \cong \mathbb{P}^5.$$

Restricting to the collection of σ-isotropic ideals, one obtains a bijection with the collection of 1-dimensional s_σ-isotropic subspaces of $\mathrm{Symd}(A,\sigma)^0$. Hence, we have an isomorphism

$$\mathrm{IV}_2(A,\sigma) \xrightarrow{\sim} \{s_\sigma = 0\} \subset \mathbb{P}(\mathrm{Symd}(A,\sigma)^0) \cong \mathbb{P}^4.$$

This realizes $\mathrm{IV}_2(A,\sigma)$ as a quadric hypersurface in $\mathbb{P}^4$ as stated in Krashen's result above (Proposition 2.9). That is, $\mathrm{IV}_2(A,\sigma)$ is obtained from $\mathrm{SB}_2(A)$ by intersecting with the hyperplane $\{\mathrm{Trd}_A = 0\}$.

Here we define the notions R-triviality and index of algebraic groups which will be used in the following sections to describe the Chernousov-Merkurjev framework for computing K_1-zero-cycles.

Definition 3.6. See [CM01]. Let G be an algebraic group over F. A point $x \in G(F)$ is called *R-trivial* if there is a rational morphism $f : \mathbb{P}^1 \dashrightarrow G$, defined at 0 and 1, and with $f(0) = 1$ and $f(1) = x$. The collection of all R-trivial elements of $G(F)$ is denoted $RG(F)$ and is a normal subgroup of $G(F)$. We denote the set of R-equivalence classes $G(F)/RG(F)$ by $G(F)/R$. If H is a normal closed subgroup of G then $RH(F)$ is a normal subgroup of $G(F)$ [CM98, Lemma 1.2].

Definition 3.7. Let G be an algebraic group and $\rho : G \to \mathbb{G}_m$ a character on G. The *index* of ρ, denoted $\mathrm{ind}(\rho)$, is the least positive integer in the image of the composition

$$G(F((t))) \xrightarrow{\rho} F((t))^\times \xrightarrow{v} \mathbb{Z}.$$

Here $F((t))$ is the field of formal Laurent series, v is the discrete valuation on $F((t))$ defined by taking the smallest power of t, and ρ denotes the induced character $\rho_{F((t))}$.

4. K_1-zero-cycles via R-equivalence and Main Result

For the proofs of the main results, we utilize the framework established in [CM01] which recovers groups of K_1-zero-cycles of homogeneous varieties in terms of algebraic groups and corresponding R-equivalence classes. Let G be an algebraic group over F. Assume $\rho : G \to \mathbb{G}_m$ is a nontrivial character of G with kernel H, and X is a smooth complete F-variety satisfying

(I) G is connected, reductive, and rational and H is smooth.
(II) For any field extension L/F, the degree homomorphism $\deg : \mathrm{CH}_0(X_L) \to \mathbb{Z}$ is injective.
(III) For any field extension L/F satisfying $X(L) \neq \emptyset$ the norm homomorphism $N_L^1 : A_0(X_L, K_1) \to L^\times$ is an isomorphism.

(IV) For any field extension L/F, $\mathrm{ind}(\rho_L) = \mathrm{ind}(X_L)$.

Theorem 4.1 ([CM01])**.** *For (G, ρ, X) as above, i.e., satisfying* (I)-(IV)*, there are canonical isomorphisms*

$$G(F)/RH(F) \simeq A_0(X, K_1)$$
$$H(F)/R \simeq \overline{A}_0(X, K_1).$$

Example 4.2 (Quadratic Spaces [CM01])**.** Let (V, q) be a non-degenerate quadratic space over F of dimension $n \geq 2$. The triple $(\mathbf{\Gamma}^+(V, q), \mathrm{Sn}, \mathrm{IV}(V, q))$ consisting of the Clifford group, the spinor norm, and the involution variety associated to (V, q) satisfies conditions (I)-(IV). Hence there are canonical isomorphisms

$$A_0(X, K_1) \simeq \Gamma^+(V, q)/R\,\mathrm{Spin}(V, q),$$
$$\overline{A}_0(X, K_1) \simeq \mathrm{Spin}(V, q)/R.$$

Quadratic spaces of dimension 5 correspond to algebras of degree 4 with symplectic involution under the exceptional identification of B_2 (which coincides with B_2' for $\mathrm{char}(F) \neq 2$) and C_2 [KMRT]. We may thus apply the above result to the case of symplectic involutions.

Theorem 4.3. *Let (A, σ) be an algebra of degree 4 with symplectic involution and let $X = \mathrm{IV}_2(A, \sigma)$. There is a canonical group isomorphism*

$$A_0(X, K_1) \simeq \mathrm{GSp}(A, \sigma)/\mathrm{Sp}(A, \sigma) = F^\times.$$

Moreover, $\overline{A}_0(X, K_1) = 0$.

Proof. Let $V = \mathrm{Symd}(A, \sigma)^0$. The variety $\mathrm{IV}_2(A, \sigma)$ is isomorphic to the quadric hypersurface $Q = \{s_\sigma = 0\} \subset \mathbb{P}(V)$ by Proposition 2.9 and Proposition 3.4. Thus, it suffices to compute K_1-zero-cycles on Q. Taking $G = \mathbf{\Gamma}^+(V, s_\sigma)$ and ρ the spinor norm, Example 4.2 yields canonical isomorphisms

$$A_0(Q, K_1) \simeq \Gamma^+(V, s_\sigma)/R\mathrm{Spin}(V, s_\sigma),$$
$$\overline{A}_0(Q, K_1) \simeq \mathrm{Spin}(V, s_\sigma)/R.$$

Since (A, σ) corresponds to (V, s_σ) under the group of equivalence of Proposition 3.3, we may write these quotients in terms of symplectic groups using the identifications $\Gamma^+(V, s_\sigma) = \mathrm{GSp}(A, \sigma)$ and $\mathrm{Spin}(V, s_\sigma) = \mathrm{Sp}(A, \sigma)$. Since $\mathbf{Sp}(A, \sigma)$ is rational [CM98, Prop. 2.4], it is R-trivial and we have $R\mathrm{Sp}(A, \sigma) = \mathrm{Sp}(A, \sigma)$ [Mer98]. Both isomorphisms claimed above then follow. Lastly, evaluating the short exact sequence of algebraic groups $1 \to \mathbf{Sp}(A, \sigma) \to \mathbf{GSp}(A, \sigma) \to \mathbb{G}_m \to 1$ on F-points gives $\mathrm{GSp}(A, \sigma)/\mathrm{Sp}(A, \sigma) = F^\times$. □

Consider the mixed involution flag variety $V_{\overline{2},4}(A, \sigma)$ defined above (Def. 2.5). Recall that the notation $\overline{2}$ indicates that we consider flags $J \subset I$ where J is σ-isotropic. We analyze the natural projection maps

$$\mathrm{IV}_2(A, \sigma) \xleftarrow{\varphi} V_{\overline{2},4}(A, \sigma) \xrightarrow{\psi} \mathrm{SB}_4(A)$$

and investigate cycles on these varieties in the following manner. We first show that φ has rational fibers, so that the groups of K_1-zero-cycles on $V_{\overline{2},4}(A,\sigma)$ and $\mathrm{IV}_2(A,\sigma)$ coincide (Lemma 4.4). We then identify the fiber of ψ over the generic point of $\mathrm{SB}_4(A)$ as the involution variety $\mathrm{IV}_2(D,\tau)$, where D is the underlying division algebra of A and τ is the restriction of σ to D, also a symplectic involution (Lemma 4.5). This reduction to a degree 4 algebra allows us to utilize Theorem 4.3.

Lemma 4.4. *Let (A,σ) be a central simple algebra of degree n and index 4 with symplectic involution. Let $X = V_{\overline{2},4}(A,\sigma)$ and $Y = \mathrm{IV}_2(A,\sigma)$. There is an isomorphism*

$$A_0(X,K_1) \simeq A_0(Y,K_1).$$

Proof. Let η be the generic point of $\mathrm{IV}_2(A,\sigma)$ with corresponding ideal $\mathfrak{n}$, which is a σ-isotropic right ideal of A of reduced dimension 2, and let $L = F(\eta)$ be its residue field. The field L satisfies $\mathrm{ind}(A_L) \mid 2$. We wish to identify the scheme-theoretic fiber $V_{\overline{2},4}(A,\sigma)_L$.

We first identify the fibers of $\psi : V_{\overline{2},4}(A,\sigma) \to \mathrm{IV}_2(A,\sigma)$ at the separable closure. In this case, $A = \mathrm{End}_{F_s}(V)$, and the map ψ is defined by taking a flag $W \subset U$ of F_s-vector spaces to W, where $\dim_{F_s} U = 4$, $\dim_{F_s} W = 2$ and W is σ_{F_s}-isotropic. The fiber over a fixed space W is the collection of 4-dimensional vector spaces U with $W \subset U \subset V$. Taking the quotient by W, we see that this fiber consists of 2-dimensional subspaces U/W of the $(n-2)$-dimensional vector space V/W. This is precisely the Grassmannian $\mathrm{Gr}(2,n-2)$. By descent, we identify the fiber as $\mathrm{SB}_2(B)$, where B is an L-algebra satisfying $\mathrm{ind}(B) \mid 2$. Hence, the variety $\mathrm{SB}_2(B)$ has an L-rational point and is therefore rational [Bla, Prop. 3]. The fiber $V_{\overline{2},4}(A,\sigma)_L$ is thus birational to $\mathbb{P}^N_L$, where $N = 2(n-4)$. It follows that $V_{\overline{2},4}(A,\sigma)$ is birational to $\mathrm{IV}_2(A,\sigma) \times \mathbb{P}^N_L$.

The group of K_1-zero-cycles is a birational invariant of smooth projective varieties [KM, Cor. RC.13], yielding an isomorphism

$$A_0(V_{\overline{2},4}(A,\sigma),K_1) \simeq A_0(\mathrm{IV}_2(A,\sigma) \times \mathbb{P}^N_L, K_1).$$

This isomorphism in conjunction with the Projective Bundle Theorem [EKM, Theorem 53.10], yields $A_0(V_{\overline{2},4}(A,\sigma),K_1) \simeq A_0(\mathrm{IV}_2(A,\sigma),K_1)$.

□

Lemma 4.5. *Let (A,σ) be a central simple algebra of index 4 with symplectic involution. Let D be the underlying division algebra of A and τ the restriction of σ to D. Let $X = V_{\overline{2},4}(A,\sigma)$ and $Y = \mathrm{IV}_2(D,\tau)$. There is an isomorphism*

$$A_0(X,K_1) \simeq A_0(Y,K_1).$$

Proof. Let η be the generic point of $\mathrm{SB}_4(A)$, with corresponding ideal $\mathfrak{n} \subset A$. The ideal $\mathfrak{n}$ is σ-regular, as the variety of σ-regular ideals of reduced dimension 4 is an open subvariety of $\mathrm{SB}_4(A)$ (see Def. 2.6). The fiber of $\psi : V_{\overline{2},4}(A,\sigma) \to \mathrm{SB}_4(A)$ over η is the variety $\mathrm{IV}_2(\mathfrak{n},\sigma)$ of σ-isotropic ideals $J \subset \mathfrak{n}$. By Theorem 2.8, $\mathrm{IV}_2(\mathfrak{n},\sigma) = \mathrm{IV}_2(D,\tau)$ for some degree 4 algebra D, Brauer-equivalent to A, with symplectic involution τ.

Thus, $A = M_n(D)$ and $\sigma|_D = \tau$. Since $\mathrm{ind}(A) = 4$, the variety $\mathrm{SB}_4(A)$ is rational, so that $V_{\overline{2},4}(A,\sigma)$ is birational to $\mathrm{IV}_2(D,\tau) \times \mathbb{P}^N$. Again, using the Projective Bundle Formula and birational invariance of $A_0(-, K_1)$, we have the desired isomorphism. □

Theorem 4.6. *Let (A,σ) be a central simple algebra over F of index 4 with symplectic involution. Let $X = \mathrm{IV}_2(A,\sigma)$ be the corresponding second generalized involution variety. There is a canonical group isomorphism $A_0(X, K_1) \simeq F^\times$. Moreover, $\overline{A}_0(X, K_1) = 0$.*

Proof. By Lemma 4.4 and Lemma 4.5, the group of K_1-zero-cycles on $\mathrm{IV}_2(A,\sigma)$ coincides with that of $\mathrm{IV}_2(D,\tau)$, where $A = M_n(D)$. Since D has degree 4, Theorem 4.3 provides the desired result. □

Remark 4.7. As the groups $\mathbf{GSp}(A,\sigma)$ and $\mathbf{Sp}(A,\sigma)$ are rational, one may look to use the Chernousov-Merkurjev framework directly to prove a similar statement for algebras of arbitrary index. Unfortunately, it is not known if the degree homomorphism is injective for algebras of index greater than 4.

References

[Bla] A. Blanchet. Function fields of generalized Brauer-Severi varieties. Comm. Algebra 19 (1991), no. 1, 97-118.

[CM98] V. Chernousov and A. S. Merkurjev. R-equivalence and special unitary groups. J. Algebra, 209: 175-198, 1998.

[CM01] V. Chernousov and A. S. Merkurjev. R-equivalence in spinor groups. J. Amer. Math. Soc. 14 (2001), no. 3, 509-534 (electronic).

[CM06] V. Chernousov and A. S. Merkurjev. Connectedness of classes of fields and zero-cycles on projective homogeneous varieties. Compos. Math. 142 (2006), no. 6, 1522-1548.

[EKM] R. Elman, N. Karpenko, and A. S. Merkurjev. *The algebraic and geometric theory of quadratic forms*. American Mathematical Society Colloquium Publications, vol. 56, American Mathematical Society, Providence, RI, 2008.

[Ful] W. Fulton. Intersection theory. Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 2. Springer-Verlag, Berlin, 1998.

[GMS] S. Garibaldi, A. S. Merkurjev, J. P. Serre. Cohomological invariants in Galois cohomology. University Lecture Series, 28. American Mathematical Society, Providence, RI, 2003.

[GS06] P. Gille and T. Szamuely. *Central simple algebras and Galois cohomology*. volume 101 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2006. 2000.

[Karp] N. Karpenko, Algebro-geometric invariants of quadratic forms, Leningrad (St. Petersburg) Math. J. 2 (1991), 119138.

[KM] N. A. Karpenko, A. S. Merkurjev. On standard norm varieties. (English, French summary) Ann. Sci. Éc. Norm. Supér. (4) 46 (2013), no. 1, 175-214, 2013.

[KMRT] Max-Albert Knus, Alexander Merkurjev, Markus Rost, and Jean-Pierre Tignol. *The book of involutions*. American Mathematical Society, Providence, RI, 1998. With a preface in French by J. Tits.

[Kra] D. Krashen. Zero cycles on homogeneous varieties. Adv. Math., 223(6):2022-2048, 2010.

[McF] P. K. McFaddin. The group of K_1-zero-cycles on the second generalized Severi-Brauer variety of an algebra of index 4. J. Algebra 479 (2017), 192-202..

[Mer95] A. S. Merkur'ev. Zero-dimensional cycles on some involution varieties. Zap. Nauchn. Sem. S.-Petersburg. Otdel. Mat. Inst. Steklov. (POMI), 227(Voprosy Teor. Predstav. Algebr i Grupp. 4): 93-105, 158, 1995. Proc. Sympos. Pure Math., 58, Part 1, Amer. Math. Soc., Providence, RI, 1995.

[Mer98] A. S. Merkurjev. K-Theory and Algebraic Groups. Progress in Mathematics, Vol. 169, Birkhäuser Verlag Basel/Switzerland, 1998.

[MS82] A.S. Merkur'ev, A.A. Suslin. K-cohomology of Severi-Brauer varieties and the norm residue homomorphism. (Russian) Izv. Akad. Nauk SSSR Ser. Mat. 46 (1982), no. 5, 1011-1046, 1135-1136.

[MS92] A. S. Merkurjev and A. A. Suslin. The group of K_1-zero-cycles on Severi-Brauer varieties. Nova J. Algebra Geom. 1 (1992), no. 3, 297-315.

[Mil] J. Milnor. Algebraic K-theory and quadratic forms. Invent. Math. 9, 318-344, 1970.

[Pan84] I. Panin, Application of K-theory in algebraic geometry, Thesis, LOMI, Leningrad (1984).

[Pan03] I. A. Panin. The equicharacteristic case of the Gersten conjecture. Translated from Tr. Mat. Inst. Steklova 241 (2003), Teor. Chisel, Algebra i Algebr. Geom., 169–178 Proc. Steklov Inst. Math. 2003, no. 2(241), 154-163.

[PSZ] V. Petrov, N. Semenov, and K. Zainoulline. Zero cycles on a twisted Cayley plane. (English summary) Canad. Math. Bull. 51, no. 1, 114-124, 2008. 1982. Studies in the History of Modern Science, 9.

[Qui] D. Quillen. Higher algebraic K-theory I. (1973), 85-147. Lecture Notes in Math., Vol. 341.

[Ros88] M. Rost. On the spinor norm and $A_0(X, K_1)$ for quadrics. Preprint (1988), `http://www.math.uiuc.edu/K-theory/0165/spinor.pdf`.

[Ros] M. Rost. Chow groups with coefficients. Doc. Math., 1:No. 16, 319-393 (electronic), 1996.

[Sri] V. Srinivas. *Algebraic K-Theory*. Reprint of the 1996 second edition. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, 2008. xviii+341, ISBN: 978-0-8176-4736-0.

[Sus] A. A. Suslin. Algebraic K-theory and the norm residue homomorphism. Current problems in mathematics, Vol. 25, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1984, pp. 115-207.

[Swa] R. G. Swan. Zero cycles on quadric hypersurfaces. Proc. Amer. Math. Soc., 107(1): 43-46, 1989.

[Tao] D. Tao. A variety associated to an algebra with involution. J. Algebra, 168(2): 479-520, 1994.

[Voe] V. Voevodsky. On motivic cohomology with Z/l-coefficients. Ann. of Math. (2) 174, no. 1, 401-438, 2011.

[Vos] V. E. Voskresenskiĭ. Algebraic tori. Nauka, Moscow, 1977 (Russian).

[Wei] C. Weibel. *The K-Book: An Introduction to Algebraic K-Theory*. Graduate Studies in Mathematics, 145. American Mathematical Society, Providence, RI, 2013.

[Weil] A. Weil, Algebras with involutions and the classical groups, J. Indian Math. Soc. 24 (1960), 589623. MR 25:147.

Department of Mathematics, University of South Carolina, Columbia, SC 29208
E-mail address: `pkmcfaddin@gmail.com`
URL: `http://mcfaddin.github.io/`